

An easy to use, oral
bronchodilator from Dechra...



DILATEROL®

25 µg/ml syrup

WHAT IS DILATEROL?

Dilaterol is a syrup containing **25 µg/ml clenbuterol hydrochloride**

- Indication:** For treatment of respiratory disease in horses where it is considered that airway obstruction due to bronchospasm and/or accumulation of mucus is a contributing factor, and improved mucociliary clearance is desirable.
- Species:** Horses
- Presentation:** 355 ml bottle containing oral syrup, with accurate pump dispenser


WHY USE DILATEROL?

- ✓ Oral clenbuterol in an easily and accurately administered liquid formulation
- ✓ Causes intense bronchodilation, inhibits histamine release and increases ciliary mucous clearance
- ✓ Particularly useful in cases of respiratory allergy, bronchospasm and equine asthma
- ✓ To be used alone or as adjuvant therapy

HOW TO USE DILATEROL?

- Administration:** Oral
The syrup should be added to the feed.
- Dose:** 0.8 µg clenbuterol hydrochloride per kg bodyweight twice daily for as long as required
- Withdrawal time:** Meat and offal – 28 days

		Dilaterol		
Species	BW (kg)	Clenbuterol hydrochloride (µg)	Volume (ml)	Pumps
	125	100	4	1
	250	200	8	2
	375	300	12	3
	500	400	16	4
	625	500	20	5



For more details, please contact us or visit our website www.dechra.co.uk/products/equine



SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Dilaterol 25 micrograms/ml syrup for horses.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ml contains:

Active substance

Clenbuterol hydrochloride 25 micrograms
(corresponding to 22 micrograms clenbuterol)

Excipients:

Methyl parahydroxybenzoate (E218) 2.02 mg
Propyl parahydroxybenzoate 0.26 mg
For the full list of excipients, see section 6.1

3. PHARMACEUTICAL FORM

Syrup
Clear colourless syrup

4. CLINICAL PARTICULARS

4.1 Target Species

Horses

4.2 Indications for use, specifying the target species

Treatment of respiratory disease in horses where it is considered that airway obstruction due to bronchospasm and/or accumulation of mucus is a contributing factor, and improved mucociliary clearance is desirable. To be used alone or as adjuvant therapy.

4.3 Contraindications

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

Do not use in horses with known cardiac disease.

For use during pregnancy or lactation see section 4.7.

4.4 Special warnings for each target species

None

4.5 Special precautions for use

Special precautions for use in animals

In cases accompanied by bacterial infection the administration of antimicrobial agents is recommended.

In case of glaucoma the product must only be used after a careful risk-benefit assessment.

Special precautions should be taken in case of halothane anaesthesia, since the heart function can show increased sensitivity to catecholamines.

Special precautions to be taken by the person administering the medicinal product to animals

This product contains clenbuterol hydrochloride, a beta-agonist.

Wear gloves to avoid skin contact. In case of accidental skin contact, wash affected area thoroughly. If irritation occurs/persists seek medical advice. Wash hands thoroughly after using the product.

Take care to avoid eye contact. In the case of accidental eye contact, flush thoroughly with clean water and seek medical advice.

Do not eat, drink or smoke when using this product. In case of accidental ingestion, seek medical advice immediately and show the package leaflet to the healthcare professional.

People with known hypersensitivity to clenbuterol should avoid contact with the veterinary medicinal product.

4.6 Adverse reactions (frequency and seriousness)

Clenbuterol may cause side effects such as sweating (mainly neck region), muscle tremor, tachycardia, slight hypotension or restlessness. These are typical for β -agonists and occur rarely.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals displaying adverse reaction(s) during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals)
- uncommon (more than 1 but less than 10 animals in 1,000 animals)
- rare (more than 1 but less than 10 animals in 10,000 animals)
- very rare (less than 1 animal in 10,000 animals, including isolated reports).

4.7 Use during pregnancy, lactation or lay

If used during pregnancy, treatment must be discontinued a minimum of 4 days before the expected time of delivery, since uterine contractions may be abolished or labour may be prolonged under its influence.

The safety of the veterinary medicinal product has not been established during lactation.

A nursing foal ingests a high volume of milk relative to its body weight. Therefore, during lactation an effect of the active substance excreted in milk in the nursing foal cannot be definitely excluded.

4.8 Interaction with other medicinal products and other forms of interaction

The product antagonises the effects of prostaglandin F₂-alpha and oxytocin.

The product is antagonised by β -adrenergic blocking agents.

Do not administer concurrently with other beta-adrenergic agents.

During the use of both local and general anaesthetics one cannot exclude a further vascular dilatation and fall of blood pressure, particularly if used in combination with atropine.

4.9 Amounts to be administered and administration route

For oral use.

Each depression of the pump delivers 4 ml of product (0.100 mg of clenbuterol hydrochloride, equivalent to 0.088 mg clenbuterol).

The pump needs to be primed before the first use only. Prime the pump by pressing twice and discard the retrieved syrup.

It is not possible to extract all the contents using the pump provided.

Administer 4 ml of the product per 125 kg bodyweight twice daily.

This is equivalent to twice daily administration of 0.8 micrograms clenbuterol hydrochloride per kg bodyweight.

The syrup should be added to the feed.

Treatment should continue for as long as necessary.

4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary

Dosages of clenbuterol hydrochloride up to 4 times the therapeutic dose

(administered orally) for a period of 90 days caused transient side effects typical for beta2-adrenoceptor agonists (sweating, tachycardia, muscle tremor), which required no treatment.

In case of accidental overdose, a β -blocker (such as propranolol) may be used as antidote.

4.11 Withdrawal period(s)

meat and offal – 28 days

Not authorized for use in lactating animals producing milk for human consumption.

5. PHARMACOLOGICAL or IMMUNOLOGICAL PROPERTIES

Pharmacotherapeutic group: selective beta-2-adrenoreceptor agonists

ATC Vet code : QR03CC13

5.1 Pharmacodynamic properties

The product contains clenbuterol hydrochloride, which is a sympathomimetic amine which preferentially binds to β_2 adrenoreceptors on cell membranes of the bronchi. This subsequently activates the enzyme adenylate cyclase in smooth muscle cells, thus providing intense bronchodilating properties and decreasing airway resistance with minimum effect on the cardiovascular system. The product has been shown to inhibit histamine release from mast cells in the lungs, and enhance mucociliary clearance in horses

5.2 Pharmacokinetic particulars

After oral administration in horses, clenbuterol is readily absorbed and maximum plasma concentrations reached within 2 hours of dosing. Steady state concentrations in plasma are reached after 3-5 days treatment and range from 1.0 – 2.2 ng/ml.

The substance is rapidly distributed in tissues and metabolised primarily by the liver. Clenbuterol is the main excretory product and approximately 45% of the dose is eliminated unchanged in the urine. The kidneys excrete 70 – 91% of the total dose, and the remainder is eliminated in the faeces (6 – 15%).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Methyl parahydroxybenzoate (E218)

Propyl parahydroxybenzoate

Carbomer 974P

Sucrose

Macrogol 400

Glycerol (85%)

Ethanol (96%)

Sodium hydroxide

Water, purified

6.2 Major incompatibilities

None known.

6.3 Shelf-life

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years

Shelf life after first opening the immediate packaging: 3 months

6.4 Special precautions for storage

Do not store above 25°C. Protect from light.

6.5 Nature and composition of immediate packaging

355 ml HDPE bottle sealed with an aluminium/PE heat seal or a transparent HDPE cap. The product is supplied in a carton box with a multi-component mechanical pump dispenser capable of delivering 4 ml of the product.

6.6 Special precautions for the disposal of unused veterinary medicinal

products or waste materials derived from the use of such products

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Le Vet Beheer B.V.

Wilgenweg 7

3421 TV Oudewater

Netherlands

8. MARKETING AUTHORISATION NUMBER(S)

Vm 41821/4001

VPA: 10475/002/001

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

7th March 2013

10. DATE OF REVISION OF THE TEXT

May 2018

Learn more online

 www.dechra.co.uk/therapy-areas/equine

Call our equine team

 01939 211200

Dechra Veterinary Products Limited, Sansaw Business Park, Hadnall, Shrewsbury, Shropshire SY4 4AS Tel: +44 (0)1939 211200
Fax: +44 (0)1939 211201 www.dechra.co.uk Registered Office: 24 Cheshire Avenue, Cheshire Business Park, Lostock Gralam, Northwich,
CW9 7UA Company Registration No.5385888. Dechra Veterinary Products Limited is a trading business of Dechra Pharmaceuticals PLC.

Dilaterol contains clenbuterol hydrochloride 25 µg/ml UK:   Sch 4 IE: 

May 2020 DVP1353


Dechra
Veterinary Products